



INFECTION

CONTROL POLICIES

& PROCEDURES

Follow us on the following UQUDENT accounts:























1.General instructions regarding environment of the clinics



CONTROL POLICIES

& PROCEDURES



The doors of the clinics should be kept closed





Food and drink are strictly prohibited inside the clinics






No Food or Drink

•All personal belongings should be left outside the clinics













INFECTION
CONTROL POLICIES
& PROCEDURES

Closed-toe shoes and socks must be worn.

No sandals, slippers, or open crocs.







For female DHCWs

1. Hair should be covered





2. Long head covers should be tightly tucked inside the clinic coat or gown





3. Clean short fingernails must be maintained





4. Nail polish, artificial nails, and nail art are not permitted




5. No hand or wrist jewelry, wristwatches, or necklaces, is accepted



For male DHCWs

1. Neckties should be tucked inside the clinic coat or gown





2. No wristwatches or other rings







Prof. Ibtesam Kamel Afifi



I. Infection Control during the pretreatment period (Before patient entry)




1. Remove unnecessary items from the dental procedure area



2. Review patient records before initiating treatment

N.B. Do not leave the record on the countertop or handle it with contaminated gloves





3. Perform hand hygiene then don a mask and gloves to prepare the clinic








4. Flush dental unit waterlines: at the beginning of the working day for 3 minutes, to prevent biofilm formation.



5. Disinfect and wrap the clinical contact surfaces of the dental chair





6. Set out all instruments, medications, impression materials, and other items that are needed for a procedure.





7. Perform Hand hygiene and dry thoroughly









8. Don personal protective equipment in the following sequence:

a) Gown







b) Mask





INFECTION
CONTROL POLICIES
& PROCEDURES

Helpful hints for mask use

- The outside surface is considered highly contaminated and should not be touched with ungloved hands.
- Masks should be changed with <u>every</u> patient or if they become wet.





Prof. Ibtesam Kamel Afifi



A damp or wet mask is *not* an effective mask.

Masks should *not* be worn under the nose or chin.



c) Protective eye wears i.e goggle or face shield







For male DHCWs

Facial hair should be completely covered under mask and face shields







d) Gloves







CORRECT SEQUENCE FOR **DONNING** PERSONAL PROTECTIVE EQUIPMENT (PPE)

The type of PPE used will vary based on the level of precautions required; e.g., Standard and Contact, Droplet or Airborne Infection Isolation.

Remove hand jewellery and tie back hair.

Clean and dry hands thoroughly.

1. GOWN / APRON

Fully cover torso from neck to knees, arms to end of wrists, and wrap around the back

Fasten in back of neck and waist



2. MASK OR RESPIRATOR

Secure ties or elastic bands at middle of head and neck

Fit flexible band to nose bridge

Fit snug to face and below chin

Fit-check respirator





3. **GOGGLES OR FACE SHIELD**

If you wear glasses put them on.

Place goggles or face shield over face and eyes and adjust to fit








4. GLOVES

Extend to cover wrist







TPOKO KI TE URU HAUDRA Infection control January 2005, Developed using CDC Guidelines 2005


DRESS CODE

Male Students



DRESS CODE

Female Students















Short Nails





Jewelries

No Jewelry





Open Toe Shoes

Follow us on the following UQUDENT accounts

























Follow us on the following UQUDENT accounts



























CONTROL POLICIES

& PROCEDURES

1. Apply dental bibs to the patient






2. Avoid touching unprotected switches, handles, and other equipment once gloves have been contaminated



3. Don't touch your mobile phone or door knobs with contaminated gloves before hand hygiene.



4. An instrument that is dropped must NOT be picked up and reused




Policy

5. Personal protective equipment should not be worn outside the clinic





6. If sharps will be used, hand- free technique for safe handling is recommended.







- 7. Take special precautions with syringes & needles
- a) Don't recap or bend the needle





b) If necessary, use one hand scoop technique




Figure 27-4 If it is necessary to a recap a needle, use a one-handed scoop technique: (A) Preparing to slide needle into cap. (B) Lifting cap onto needle. (C) Covering needle with cap. Alternatively, a needle-recapping device may be used.




c) Dispose used needles in sharps box





Be careful on returning hand pieces into their tie in sockets







- 7. Immediately manage exposure to blood by sharps injury, needle stick, or splash:
 - a) Wash wound / exposed area with soap and water.
 - b) Irrigate eyes with sterile water if availablec)cover the wound with a waterproof plaster or dressing



الاسعافات الأولية عند الإصابة بالألات الحادة أو وخز الإبر First Aid after Needle Stick or Sharp Object Injury



توقف عن العمل Stop working



2 شجع النزيف الحر Encourage free bleeding



3 نظف الجرح برفق بالماء والصابون Cleanse the wound with soap and water



جفف الجرح وقم بتغطيته بالبلاستر Dry and cover the wound with plaster

عند الإصابة بالألات الحادة أو وخرز الإبر اتبع In case of Needle Stick or Sharp Object Injury perform the following steps:

- 1. Stop working.
- 2. Encourage free bleeding gently by running cool water over the bleeding area for several minutes.
- 3. Do NOT squeeze the wound to encourage bleeding.
- 4. Gently cleanse the wound with soap and water (Do not scrub the wound while you're washing it).
- 5. Dry and cover the wound with a waterproof plaster or dressing.
- 6. Notify infection control supervisor about the injury.
- 7. Fulfil needle stick and sharp object injury report form with the help of infection control supervisor.
- 8. Perform laboratory testing and medical follow-up according to the given advice.

الخطوات التالية:

- ١. توقف عن العمل.
- ٢. شجع النزيف الحر بلطف عن طريق تشغيل ماء بارد على منطقة النزيف لعدة دقائق.
 - ٣. لا تضغط على الجرح لايقاف النزيف.
 - ٤. نظف الجرح برفق بالماء والصابون بدون دعك.
 - ٥. جفف الجرح وقم بتغطيته بالبلاستر.
 - ٦. بلغ مشرف مكافحة العدوى عن الإصابة.
- ٧. قم بملئ نموذج الإبلاغ عن الإصابة بالألات الحادة أو وخز الإبر بمساعدة طبيب مكافحة العدوى.
- ٨. اجرى الاختبارات المعملية مع المتابعة الطبية وفقاً للنصيحة المقدمة.

d) Report to ICC and fill sharps injury form





	did the injury	, occur.	
	Before use of item (item broke / slipped, assembling device).		
	During use of item (item slipped, patient jarred item).		
	Restraining patient		
	o instr	uments).	tep procedure (between incremental injections, passing
	Disassembling device or equipment.		
	In preparation for reuse of reusable instrument (sorting, disinfecting, sterilizing).		
	While recapping used needle. Device left on the floor, table or other inappropriate place.		
	After use-before disposal.		
	While putting item into disposable container.		
0	After disposal, stuck by item protruding from opening of disposal container.		
	Item pierced side of disposal container.		
	After disposal, item protruded from trash bag or inappropriate waste container.		
	Other, descr	iption of incident	t: 🔛
THE LO	CATION OF T	HE INJURY	
EXPO	SED PATIENT:		
Did th	e exposed per	rson vaccinated ag	against HBV?
		□Yes	□No
Did th	e exposed per	son detect his imn	nmune state before?
		☐ Yes	□No
If yes,	what is his im	mune state?	
		Responder	☐ Non responder
SOUR	E PATIENT:		
Does s	ero status of t	he source patient	t is known regarding:
HBs A	3	☐ Positive	□ Negative
HCV a	ntibodies	☐ Positive	Negative
HIV an	tibodies	☐ Positive	□ Negative
ICC me	ember comple	ting the report:	



9. If you perform x-ray to the patient, you should:

- a) Disinfect and wrap x-ray handle, button, and computer mouse
- b) Perform hand hygiene, dry your hand, and wear new gloves









CONTROL POLICIES

& PROCEDURES

- 1. Begin the cleaning and disinfection process by:
- a) Removing contaminated gloves used during treatment.
- b) Perform hand hygiene
- c) Put on another pair of gloves before beginning the clean-up. Continue to wear protective eyewear, mask, and gown.





- 2. Waste disposal:
- a) Discard used disposable items immediately.
- b) Remove the tray with all instruments and deliver it to the dental assistant to provide it to the sterilization/clean-up area.
- c) Dispose extracted teeth in clinical waste containers



3. Remove all disposable barriers





4. Clean and disinfect all items including those not protected by barriers.







5. Remove PPE in the following sequence:

CORRECT SEQUENCE FOR REMOVING PERSONAL PROTECTIVE EQUIPMENT (PPE) 1. GLOVES Outside of gloves are contaminated—DO NOT TOUCH! Grasp outside of glove with opposite glovedhand; peel off Hold removed glove in gloved hand Slide fingers of ungloved hand under remaining glove at wrist



Peel glove off over first glove

Discard gloves in waste container

Clean and dry your hands thoroughly

Outside of goggles or face shield are contaminated—DO NOT TOUCH!

To remove, handle by head band or ear pieces

Place in designated receptacle for reprocessing or in waste container

Clean and dry your hands thoroughly



Gown front and sleeves are contaminated—DO NOT TOUCH!

Unfasten ties

Pull away from neck and shoulders, touching inside of gown only

Turn gown inside out

Fold or roll into a bundle and discard

Clean and dry your hands thoroughly



Front of mask/respirator is contaminated—DO NOT TOUCH!

Grasp bottom, then top ties or elastics and remove

Discard in waste container

Clean and dry your hands thoroughly







6.Perform Hand hygiene and dry thoroughly









Refer to UQUDENT manual of infection control policies & Procedures

