Lecture 5





Identification of bacteria

Objectives

- Define bacterial colony and describe how to obtain pure culture
- Enumerates basic categories of laboratory tests to identify bacteria [colony morphology, microscopy, single enzyme tests, Multi enzyme tests, molecular methods, and immunologic methods].
- Describe the principal and describe and example of each category of laboratory tests.
- Compare direct and indirect ELISA techniques.


Colony


A <u>colony</u> is visible mass of large number of microorganisms, growing on or in solid medium.



Pure Culture


It is essential to start <u>identification</u> tests with <u>pure bacterial</u> <u>isolates</u> grown from a single colony.





A. Colony morphology

- A <u>colony</u> is a pile or mass of a sufficiently large number of cells, growing on or in solid medium, that they are visible to the naked eye.
- Different species of bacteria can produce very different colonies.



1. Colony morphology

Example of variation in colony morphology include:

- Colony size,
- Colony Edge/margin
- Colony color

- Colony shape,
- Colony Elevation
- Presence of hemolysis on blood agar


etc.....


1. Colony morphology



Colony Shape

1. Colony morphology







The shape of the edge or margin can be determined by looking down at the top of the colony.

The nature of the colony elevation is apparent when viewed from the side as the plate is held at eye level.







Types of Hemolysis on Blood agar



B. Microscopy and Gram stain

To determine bacterial morphology and Gram reaction.

Bacterial Morphology Include:

- Size • Shape
- Spore
- Staining

- Capsule
- Motility

• Arrangement

C. Single-enzyme tests

- Different bacteria produce varying spectra of enzymes.
- Example → Catalase test:


Catalase test

• The enzyme catalase catalyzes the degradation of hydrogen peroxide to water and molecular oxygen

 $(H_2O_2 \rightarrow H_2O + O_2).$

Catalase test

Catalase-positive organisms rapidly produce bubbles when exposed to a solution containing hydrogen peroxide.




Catalase positive test Catalase negative test D. Tests based on the presence of metabolic pathways

- These tests measure the presence of a metabolic pathway in a bacterial isolate, rather than a single enzyme.
- Example → Sugar fermentation tests.

Sugar fermentation tests:

To determine the ability of the bacteria to ferment a particular sugar the bacteria is inoculated in a test tube containing:


- The test sugar solution
- A pH indicator which change its color if acid is produced.
- An **inverted tube** which is used to detect gas production.



Sugar Fermentation tests

There are three possible results

- That bacteria can't ferment the test sugar (1)
- That bacteria can ferment the test sugar with gas production (2)
- That bacteria can ferment the test sugar but without gas production (3)



Uninoculated API strip



E. Automated systems

Microbiology laboratories are increasingly using automated methods to identify bacterial pathogens.



F. Immunologic Detection of Microorganisms

Immunologic techniques can be used for identification of microorganisms by:


1. Detection of bacterial **antigens** in specimens or cultures: using known specific antibody.

2. Detection in a patient's serum of **antibodies** that are directed against microbial antigens (Serology).

Immunologic Detection of Microorganisms

The presence of **antibodies** or **antigens** in patients' sera or other body fluids is determined by performing an **immunoassay procedure**, such as:


- Slide Agglutination.
- Enzyme-linked immunosorbent assay (ELISA).



- Detection of <u>bacterial antigens</u> provides a direct evidence that the patient is currently infected.
- Detecting **antibodies** to a particular pathogen may represent:
 - present infection,
 - **2** past infection, or
 - prior vaccination against that pathogen.


Agglutination test

Agglutination is the visible clumping of particulate (insoluble) antigen e.g. bacteria or red blood cells with its specific antibody.



Slide agglutination test

- Agglutination reactions can be used for rapid identification of bacteria (slide agglutination test).
- A drop of bacterial suspension is mixed with specific antibacterial antibody on a microscopic slide.
- This test is repeated with different antibodies to detect which antibody causes agglutination.



- Enzyme-linked immunosorbent assay (ELISA) is a sensitive assay useful in detecting either antigens or antibodies in <u>low concentrations</u> in a patient's body fluids.
- There are two types of ELSIA technique:
 - **<u>Direct ELISA</u>** technique is used to detect **<u>antigen</u>** in specimens.
 - Indirect ELISA to detect and measure antibodies in serum.




Microtitration plate Positive Test is indicated by color change


Direct ELISA (Sandwich method)

For ANTIGEN detection

For ANTIBODY detection





Well of microtitration plate is coated with specific known antibody




Indirect ELISA

Well of microtitration plate is coated with known antigen.







4



Microtitration plate Positive Test is indicated by color change

G. Molecular Methods

- The main objective of the molecular methods of bacterial identification is direct recognition of pathogen-specific nucleic acid sequences in the test material.
- These methods are used in particular in the search for bacteria that are very **difficult to culture**, or proliferate very slowly.


Molecular Methods

The following methods are used:

- DNA Probes
- Nucleic Acid Amplification


DNA probes.

- DNA probes are:
 - small segments of complementary DNA
 - labeled with a **marker** molecule (such as enzymes or radioactive label).




DNA probes.

- The DNA probes hybridizes (attaching to) its complementary nucleic acid sequences in the test sample.
- The presence and the quantity of hybrids is determined by the detection of the label.



Nucleic Acid Amplification:

The polymerase chain reaction (PCR) is a scientific technique to <u>amplify</u> a specific sequence in single or a few copies of DNA strands to generate thousands to millions of copies.









Nucleic acid amplification can be done by:

- A. Polymerase chain reaction (PCR)
- B. DNA probes
- C. Enzyme-linked immunosorbent assay (ELISA)
- D. API-20E





2. Detection of bacterial antigen in patient sample indicates _____?

- A. Present infection
- B. Past infection
- C. Previous immunization
- D. All of the above





3- Study the following diagram and then fill in the spaces:

- This test is:
- The inverted tube is used to detect:
- The change of the color of the tube indicate




4- Study the following diagram and then fill in the spaces:

A. The name of the test is:

................


- B. The tube contain a solution of:
- C. Positive test is indicated by:
- D. Tube b is (positive/negative)







Study the diagram and answer:



- 5. The name of the above test is:
- A. Polymerase chain reaction (PCR)
- B. DNA probes
- C. Enzyme-linked immunosorbent assay (ELISA)
- D. API20E

6. This test is used to detect :

- A. Bacterial Nucleic acid
- **B.** Bacterial Antigens
- C. Bacterial Morphology
- D. Bacterial Metabolic pathways





7. A visible mass of large number of microorganisms, growing on or in solid medium is called _____

- A. Colony
- B. Mold
- C. Biofilm
- D. Probe





8. A small segments of complementary DNA labeled with a marker molecule is called:

- A. Plasmid
- B. Antigen
- C. Probe
- D. mRNA
- E. Antibody




9. Complete what abbreviations stand for:

- A. ELISA:
- B. PCR:

10. Study the diagram and fill in the spaces:

- A. The Name of the above test is:
- B. The test is used for detection of:
- C. The wells are coated with:
- D. Is the above test direct or indirect:

.



11. Study the diagram and fill in the spaces:

- B. The wells are coated with: _____(Antigen/Antibody)
- C. The test is used for detection of: _____ (Antigen/ Antibody)
- D. Is the above test _____ (direct / indirect)
- E. In positive tests there will be _____
 (color change/ gas bubbles/ precipitation / agglutination).

